

QUICK REFERENCE INSTRUCTIONS

2019-nCoV Ag Rapid Test Kit

(Immunochromatography)

SAMPLE REQUIREMENTS


Insert a sterile swab into the nostril of the patient, reaching the surface of the posterior nasopharynx.



2 Swab over the surface of the posterior nasopharynx. Hold the swab in the nosefor 15-30s and rotate it 3 times.




Withdraw the sterile swab from the nasal cavity.



*Note:

- The samples should be used as soon as possible after collected (within half an hour).
- 2. Samples should not be inactivated.

SAMPLE PROCESSING




- 1 300μL (to the 300L mark line,9 drops)Sample Buffer.
- Vigorously mix at least 10 times.
- 3 Squeeze liquid from swab.
- Cover the dripper head.

TEST PROCEDURE



3 drops to the each well
NOTE:Take the required reagents and test
cards to equilibrate to room temperature.

INTERPRETATION OF TEST RESULTS



Read the result after 15mins. less than 15 minutes may lead a false negative result; more than 15 minutes may lead a false positive result.

IMPORTANT

- 1. Read instructions carefully before beginning.
- 2. All specimens must be at room temperature before testing.
- 3. Check expiration date on each individual test package or outer box before using.
- 4. Do not use any test past the expiration date on the label.
- 5. Disposable gloves and safety glasses are recommended when running this test.
- 6. Wash hands thoroughly after handling any patient sample.

Huachenyang (Shenzhen) Technology Co., Ltd
Area A, F8/Blcg 4, Hengchangrong High Tech Park, Shangnan East
Road, Hongtian, Shajing, Bao'an, Shenzhen, 518125, Guangdong, CHINA